

CALIFORNIA WILDLIFE HABITAT RELATIONSHIPS SYSTEM
maintained by the
CALIFORNIA DEPARTMENT OF FISH AND GAME
and supported by the
CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP
Database Version 8.1 (2005)

R060 Long-nosed Snake *Rhinocheilus lecontei*
Family: Colubridae Order: Squamata Class: Reptilia

Written by: S. Morey
Reviewed by: T. Papenfuss
Edited by: R. Duke

DISTRIBUTION, ABUNDANCE, AND SEASONALITY

Common only in desert regions and locally in coastal southern California, the long-nosed snake is uncommon in the interior Coast Ranges and the adjacent floor of the Central Valley from Sutter and Butte cos. south. This species ranges widely in southern California and is known from desert regions east of the Sierra and Cascade Ranges north to the Oregon border. A snake primarily of grasslands, arid brushlands, and coastal chaparral, the long-nosed snake is not expected in forested types. Elevation sea level to 1620 m (5400 ft).

SPECIFIC HABITAT REQUIREMENTS

Feeding: Long-nosed snakes prey heavily on lizards but also take rodents and other small prey. Stebbins (1954) listed mammals, lizards and lizard eggs, and insects as among the food items taken. Captive individuals have also eaten small snakes.

Cover: Seldom found under surface objects, long-nosed snakes are good burrowers. Nussbaum et al. (1983) report that disturbed individuals burrow rapidly into loose soil, and that they often lie covered with sand with only their head exposed. They may also seek cover in rock crevices (Stebbins 1954).

Reproduction: Little is known about habitat requirements for reproduction in this species. Females are thought to deposit egg clutches in loose, moist, well-aerated soil. Abandoned mammal burrows may also be utilized as nest sites.

Water: No information on water requirements. This is a snake of arid lands, seldom encountered near stream courses. Standing water is not an important habitat element.

Pattern: Inhabits deserts, plains and brushland, seldom found near water.

SPECIES LIFE HISTORY

Activity Patterns: Nocturnal and crepuscular, occasional diurnal activity has been reported. Individuals are most commonly encountered from April thru June (Klauber 1941). A period of winter inactivity occurs at all localities.

Seasonal Movements/Migration: Predictable seasonal movements have not been reported for this species in California.

Home Range: The nature of the home range in this species is unknown.

Territory: No evidence for the territorial defense of resources has been reported.

Reproduction: Five to 8 eggs are deposited in July. Hatchlings emerge in late August or September.

Niche: Long-nosed snakes appear in the diet of hawks (Nussbaum et al. 1983), and they are taken as prey by other snakes (Cunningham 1959). Competitive interactions between this somewhat fossorial species and other snakes (some of them also good burrowers) with which they co-occur are largely unknown.

REFERENCES

- Cunningham, J. D. 1959. Reproduction and food of some California snakes. *Herpetologica* 15:17-19.
- Diller, L. V., and R. C. Wallace. 1981. Additional distribution records and abundance of three species of snakes in southwestern Idaho. *Great Basin Nat.* 41:154-157.
- Klauber, L. M. 1941. The long-nosed snakes of the genus *Rhinocheilus*. *Trans. San Diego Soc. Nat. Hist.* 9:289-332.
- Medica, P. A. 1975. *Rhinocheilus*, *R. lecontei*. *Cat. Am. Amphibians and Reptiles*. 175.1-174.4.
- Nussbaum, R. A., E. D. Brodie, Jr., and R. M. Storm. 1983. *Amphibians and reptiles of the Pacific Northwest*. Univ. Press of Idaho. 332pp.
- Stebbins, R. C. 1954. *Amphibians and reptiles of western North America*. McGraw-Hill, New York. 536pp.